



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 38/43, 38/45, 39/02, C07K 16/12, 16/40, C12N 1/21, 5/10, 9/10, 15/54, 15/63	A1	(11) International Publication Number: WO 99/26651 (43) International Publication Date: 3 June 1999 (03.06.99)
(21) International Application Number: PCT/US98/25073 (22) International Filing Date: 24 November 1998 (24.11.98) (30) Priority Data: 08/978,454 25 November 1997 (25.11.97) US (71) Applicant: SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). (72) Inventors: PALMER, Leslie, M.; 2820 Eagleville Road, Audubon, PA 19407 (US). FEDON, Jason, C.; Apartment 2, 106 Fairfield Lane, Strafford, PA 19087 (US). WARREN, Richard, L.; 825 Cathcart Road, Blue Bell, PA 19422 (US). KOSMATKA, Anna, L.; 482 Pebble Hill Road, Doylestown, PA 18901 (US). SHILLING, Lisa, K.; 6 East Park Road, Newtown, PA 18940 (US). BLACK, Michael, T.; 502 Milhouse Way, Chester Springs, PA 19425 (US). HODGSON, John, E.; 260 Lapp Road, Malvern, PA 19355 (US). NICHOLAS, Richard, O.; 355 Carmen Drive, Collegeville, PA 19426 (US). KNOWLES, David, J., C.; Ladywell House, York Road, Boroughbridge, North Yorkshire (GB). LONETTO, Michael, A.; 18 Victoria Circle, Collegeville, PA 19426 (US). STODOLA, Robert, K.; 309 Haws Lane, Flourtown, PA 19031 (US).	(74) Agent: JACKSON, Arthur, E.; Dechert Price & Rhoads, 400 Bell Atlantic Tower, 1717 Arch Street, Philadelphia, PA 19103 (US). (81) Designated States: JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: ribH (57) Abstract The invention provides ribH polypeptides and polynucleotides encoding ribH polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are methods for utilizing ribH polypeptides to screen for antibacterial compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

ribH

RELATED APPLICATIONS

- 5 This application claims benefit of US Patent Application Number 08/978,454 filed November 25, 1997.

FIELD OF THE INVENTION

- 10 This invention relates to newly identified polynucleotides and polypeptides, and their production and uses, as well as their variants, agonists and antagonists, and their uses. In particular, the invention relates to novel polynucleotides and polypeptides of the Lumazine synthase (β -subunit) family, hereinafter referred to as "ribH".

BACKGROUND OF THE INVENTION

- 15 The Streptococci make up a medically important genera of microbes known to cause several types of disease in humans, including, for example, otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid. Since its isolation more than 100
20 years ago, *Streptococcus pneumoniae* has been one of the more intensively studied microbes. For example, much of our early understanding that DNA is, in fact, the genetic material was predicated on the work of Griffith and of Avery, Macleod and McCarty using this microbe. Despite the vast amount of research with *Streptococcus pneumoniae*, many questions concerning the virulence of this microbe remain. It is particularly preferred to employ Streptococcal genes and gene products as
25 targets for the development of antibiotics.

- Riboflavin (vitamin B2) is a member of the B complex of vitamins which function as coenzymes in metabolic reactions. Riboflavin has two coenzyme forms, flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) which act in oxidation-reduction reactions such as the cytochrome system of electron transport and the oxidative degradation of pyruvate, fatty
30 acids and amino acids. The first committed step in the biosynthesis of riboflavin is the opening of the imidazole ring of GTP. In the presence of 3 H₂O and Mg⁺⁺, the C-8 of GTP is released as formate accompanied by the release of pyrophosphate by the action of GTP cyclohydrolase II

(GCH2; EC 3.5.4.25). This enzyme function is encoded by *ribA* in bacteria and *rib1* in yeast species. Through a series of steps, involving 3,4-dihydroxy-2-butanone 4 phosphate synthase (*ribA*), 6,7-dimethyl-8-ribityllumazine synthase (*ribH*), riboflavin synthase (*ribB*), pyrimidine deaminase and pyrimidine reductase (*ribG*), enzymes encoded by genes within the riboflavin biosynthesis operon, riboflavin is formed. Because the genes required for riboflavin biosynthesis are present in many pathogenic microorganisms, and since riboflavin biosynthesis has shown to be required for virulence in the swine pathogen *Actinobacillus pleuropneumoniae* (Fuller, TE, *et al.* (1996) A riboflavin auxotroph of *Actinobacillus pleuropneumoniae* is attenuated in swine. Infect. Immun. 64:4659-4664), these gene products represent broad spectrum antibacterial as well as antifungal targets.

The frequency of *Streptococcus pneumoniae* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Streptococcus pneumoniae* strains which are resistant to some or all of the standard antibiotics. This phenomenon has created a demand for both new anti-microbial agents, vaccines, and diagnostic tests for this organism.

Clearly, there exists a need for factors, such as the *ribH* embodiments of the invention, that have a present benefit of being useful to screen compounds for antibiotic activity. Such factors are also useful to determine their role in pathogenesis of infection, dysfunction and disease. There is also a need for identification and characterization of such factors and their antagonists and agonists to find ways to prevent, ameliorate or correct such infection, dysfunction and disease.

Certain of the polypeptides of the invention possess amino acid sequence homology to a known *Actinobacillus pleuropneumoniae* *ribH* protein. (Swiss Prot. Accession no. P50856; GenBank Accession no. U27202; Perkins, J.B. *et al.*, In: *Bacillus subtilis and Other Gram-Positive Bacteria*. Eds: Sonenshein, A.L. *et al.*, 1993. American Society for Microbiology.)

SUMMARY OF THE INVENTION

It is an object of the invention to provide polypeptides that have been identified as novel *ribH* polypeptides by homology between the amino acid sequence set out in Table 1 [SEQ ID NO: 2 or 4] and a known amino acid sequence or sequences of other proteins such as *Actinobacillus pleuropneumoniae* *ribH* protein.

It is a further object of the invention to provide polynucleotides that encode ribH polypeptides, particularly polynucleotides that encode the polypeptide herein designated ribH.

In a particularly preferred embodiment of the invention, the polynucleotide comprises a region encoding ribH polypeptides comprising a sequence set out in Table 1 [SEQ ID NO:1 or 3]
5 which includes a full length gene, or a variant thereof.

In another particularly preferred embodiment of the invention, there is a novel ribH protein from *Streptococcus pneumoniae* comprising the amino acid sequence of Table 1 [SEQ ID NO:2 or 4], or a variant thereof.

As a further aspect of the invention, there are provided isolated nucleic acid molecules
10 encoding ribH, particularly *Streptococcus pneumoniae* ribH, including mRNAs, cDNAs, genomic DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

In accordance with another aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic
15 immunization. Among the particularly preferred embodiments of the invention are naturally occurring allelic variants of ribH and polypeptides encoded thereby.

As another aspect of the invention, there are provided novel polypeptides of *Streptococcus pneumoniae* referred to herein as ribH as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

20 Among the particularly preferred embodiments of the invention are variants of ribH polypeptide encoded by naturally occurring alleles of the ribH gene.

In a preferred embodiment of the invention, there are provided methods for producing the aforementioned ribH polypeptides.

In accordance with yet another aspect of the invention, there are provided inhibitors to
25 such polypeptides, useful as antibacterial agents, including, for example, antibodies.

In accordance with certain preferred embodiments of the invention, there are provided products, compositions and methods for assessing ribH expression, treating disease, assaying genetic variation, and administering a ribH polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Streptococcus pneumoniae* bacteria.

30 In accordance with certain preferred embodiments of this and other aspects of the invention, there are provided polynucleotides that hybridize to ribH polynucleotide sequences, particularly under stringent conditions.

In certain preferred embodiments of the invention, there are provided antibodies against ribH polypeptides.

In other embodiments of the invention, there are provided methods for identifying compounds which bind to or otherwise interact with and inhibit or activate an activity of a polypeptide or polynucleotide of the invention comprising: contacting a polypeptide or polynucleotide of the invention with a compound to be screened under conditions to permit binding to or other interaction between the compound and the polypeptide or polynucleotide to assess the binding to or other interaction with the compound, such binding or interaction being associated with a second component capable of providing a detectable signal in response to the binding or interaction of the polypeptide or polynucleotide with the compound; and determining whether the compound binds to or otherwise interacts with and activates or inhibits an activity of the polypeptide or polynucleotide by detecting the presence or absence of a signal generated from the binding or interaction of the compound with the polypeptide or polynucleotide.

In accordance with yet another aspect of the invention, there are provided ribH agonists and antagonists, preferably bacteriostatic or bacteriocidal agonists and antagonists.

In a further aspect of the invention, there are provided compositions comprising a ribH polynucleotide or a ribH polypeptide for administration to a cell or to a multicellular organism.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to novel ribH polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel ribH of *Streptococcus pneumoniae*, which is related by amino acid sequence homology to *Actinobacillus pleuropneumoniae* ribH polypeptide. The invention relates especially to ribH having the nucleotide and amino acid sequences set out in Table 1 as SEQ ID NO: 1 and SEQ ID NO: 2, respectively.

TABLE 1

ribH Polynucleotide and Polypeptide Sequences

- 5 (A) Sequences from *Streptococcus pneumoniae* ribH polynucleotide sequence [SEQ ID NO:1].

5' -

ATGAACACTTATGAAGGTAATTTAGTAGCAAACAATATTTAAATAGGTATTGTTGTAGCGAGATTTAAT
 GAATTTATAACTTCAAATTATTATCTGGAGCACTAGATAATCTCAAAAGAGAGAATGTAAACGAGAAA
 10 GATATCGAGGTAGCCTGGGTCCAGGAGCTTTTGAAATACCACTGATTGCATCAAAAATGGCAAAAAGT
 AAAAAATATGATGCAATTATCTGCTTGGGAGCTGTCATTAGAGGGAATACAAGTCATTATGATTATGTA
 TGTCAGCGAGGTATCTAAAGGAATCGCCCAAATCAGTTTAAATAGCGAAATTCCTGTTATGTTTGGTGTG
 CTAACGACAGATAACAATTGAACAAGCCATAGAACGAGCTGGTACTAAAGCAGGAAATAAGGGTTCTGAG
 TGTGCACAAGGAGCTATTGAAATGGTCAACCTAATTCGTACATTAGACGCATAG-3'

15

- (B) *Streptococcus pneumoniae* ribH polypeptide sequence deduced from the polynucleotide sequence in this table [SEQ ID NO:2].

NH₂-

MNTYEGNLVANNIKIGIVVARFNEFITSKLLSGALDNLKRENVNEKDIEVAWVPGAFEIPLIASKMAKS
 20 KKYDAIICLGAVIRGNTSHYDYVCSEVSKGIAQISLNSEIPVMFGVLTTDTIEQAIERAGTKAGNKGSE
 CAQGAIEMVNLIRTLDA-COOH

- (C) Polynucleotide sequences comprising *Streptococcus pneumoniae* ribH ORF sequence [SEQ ID NO:3].

25 5' -

CTAGTTTGCTCTATGATTTTCACAGAGTATTAAATTGCGATTTTGCCAAGTTTCTTTATTTCGTCTAAAA
 GTAGAGTCTGTTCTATGCGTCTAATGTACGAATTAGGTTGACCATTTCAATAGCTCCTTGTGCACACTC
 AGAACCCCTTATTTCTGCTTTAGTACCAGCTCGTTCTATGGCTTGTTCAATTGTATCTGTCGTTAGCAC
 ACCAAACATAACAGGAATTTGCTATTTAAACTGATTTGGGCGATTCCTTTAAATACCTCGCTACATAC
 30 ATAATCATAATGACTTGTATTCCCTCTAATGACAGCTCCCAAGCAAATAATTGCATCATATTTTTTACT
 TTTTGCCATTTTTCATGCAATCAGTGGGTATTTCAAAA
 -3'

(D) *Streptococcus pneumoniae* ribH polypeptide sequence deduced from the polynucleotide ORF sequence in this table [SEQ ID NO:4].

NH₂-

LKYPLIASKMAKSKKYDAIICLGAVIRGNTSHYDYVCSEVFKGIAQISLNSEIPVMFGVLTTDTIEQAI
5 ERAGTKAGNKGSECAQGAITEMVNLIRTLDA-COOH

Deposited materials

A deposit containing a *Streptococcus pneumoniae* 0100993 strain has been deposited with the National Collections of Industrial and Marine Bacteria Ltd. (herein "NCIMB"), 23 St. Machar
10 Drive, Aberdeen AB2 1RY, Scotland on 11 April 1996 and assigned deposit number 40794. The deposit was described as *Streptococcus pneumoniae* 0100993 on deposit. On 17 April 1996 a *Streptococcus pneumoniae* 0100993 DNA library in *E. coli* was similarly deposited with the NCIMB and assigned deposit number 40800.. The *Streptococcus pneumoniae* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

15 The deposited strain contains the full length ribH gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent
20 Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.


A license may be required to make, use or sell the deposited strain, and compounds derived
25 therefrom, and no such license is hereby granted.

One aspect of the invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Streptococcus pneumoniae* 0100993 strain contained in the deposited strain. Further provided by the invention are ribH nucleotide sequences of the DNA in the deposited strain and amino acid sequences encoded thereby. Also provided by the invention are ribH
30 polypeptide sequences isolated from the deposited strain and amino acid sequences derived therefrom.

Polypeptides

The polypeptides of the invention include a polypeptide of Table 1 [SEQ ID NO:2 or 4] (in particular the mature polypeptide) as well as polypeptides and fragments, particularly those which have the biological activity of ribH, and also those which have at least 70% identity to a polypeptide of Table 1 [SEQ ID NO:1 or 3] or the relevant portion, preferably at least 80% identity to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and more preferably at least 90% similarity (more preferably at least 90% identity) to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to a polypeptide of Table 1 [SEQ ID NO:2 or 4] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

The invention also includes polypeptides of the formula:



wherein, at the amino terminus, X is hydrogen, and at the carboxyl terminus, Y is hydrogen or a metal, R_1 and R_3 are any amino acid residue, m is an integer between 1 and 1000 or zero, n is an integer between 1 and 1000 or zero, and R_2 is an amino acid sequence of the invention, particularly an amino acid sequence selected from Table 1. In the formula above R_2 is oriented so that its amino terminal residue is at the left, bound to R_1 , and its carboxy terminal residue is at the right, bound to R_3 . Any stretch of amino acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer.

A fragment is a variant polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned polypeptides. As with ribH polypeptides fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region, a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of Table 1 [SEQ ID NO:2 or 4], or of variants thereof, such as a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus. Degradation forms of the polypeptides of the invention in a host cell, particularly a *Streptococcus pneumoniae*, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic

regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Also preferred are biologically active fragments which are those fragments that mediate activities of ribH, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those fragments that are antigenic or immunogenic in an animal, especially in a human. Particularly preferred are fragments comprising receptors or domains of enzymes that confer a function essential for viability of *Streptococcus pneumoniae* or the ability to initiate, or maintain cause disease in an individual, particularly a human.

Variants that are fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention.

In addition to the standard single and triple letter representations for amino acids, the term "X" or "Xaa" may also be used in describing certain polypeptides of the invention. "X" and "Xaa" mean that any of the twenty naturally occurring amino acids may appear at such a designated position in the polypeptide sequence.

Polynucleotides

Another aspect of the invention relates to isolated polynucleotides, including the full length gene, that encode the ribH polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2 or 4] and polynucleotides closely related thereto and variants thereof.

Using the information provided herein, such as a polynucleotide sequence set out in Table 1 [SEQ ID NO:1 or 3], a polynucleotide of the invention encoding ribH polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Streptococcus pneumoniae* 0100993 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a sequence given in Table 1 [SEQ ID NO:1 or 3], typically a library of clones of chromosomal DNA of *Streptococcus pneumoniae* 0100993 in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent conditions. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently, such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone.

Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Table 1 [SEQ ID NO:1 or 3] was discovered in a DNA library derived from *Streptococcus pneumoniae* 0100993.


The DNA sequence set out in Table 1 [SEQ ID NO:1 or 3] contains an open reading frame encoding a protein having about the number of amino acid residues set forth in Table 1 [SEQ ID NO:2 or 4] with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known in the art. The polynucleotide of SEQ ID NO: 1, between nucleotide number 1 and the stop codon which begins at nucleotide number 466 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

The ribH polypeptide of the invention is structurally related to other proteins of the Lumazine synthase (β -subunit) family.

The invention provides a polynucleotide sequence identical over its entire length to a coding sequence in Table 1 [SEQ ID NO:1 or 3]. Also provided by the invention is the coding sequence for the mature polypeptide or a fragment thereof, by itself as well as the coding sequence for the mature polypeptide or a fragment in reading frame with other coding sequence, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence. The polynucleotide may also contain non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences, termination signals, ribosome binding sites, sequences that stabilize mRNA, introns, polyadenylation signals, and additional coding sequence which encode additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA tag (Wilson *et al.*, *Cell* 37: 767 (1984). Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

A preferred embodiment of the invention is a polynucleotide of comprising nucleotide 1 to the nucleotide immediately upstream of or including nucleotide 466 set forth in SEQ ID NO:1 of Table 1, both of which encode the ribH polypeptide.

The invention also includes polynucleotides of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R_1 and R_3 is any nucleic acid residue, m is an integer between 1 and 3000 or zero, n is an integer between 1 and 3000 or zero, and R_2 is a nucleic acid sequence of the invention, particularly a nucleic acid sequence selected from Table 1. In the polynucleotide formula above R_2 is oriented so that its 5' end residue is at the left, bound to R_1 , and its 3' end residue is at the right, bound to R_3 . Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a heteropolymer or a homopolymer, preferably a heteropolymer. In a preferred embodiment m and/or n is an integer between 1 and 1000.

It is most preferred that the polynucleotides of the inventions are derived from *Streptococcus pneumoniae*, however, they may preferably be obtained from organisms of the same taxonomic genus. They may also be obtained, for example, from organisms of the same taxonomic family or order.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Streptococcus pneumoniae* ribH having an amino acid sequence set out in Table 1 [SEQ ID NO:2 or 4]. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or an insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode for variants of the polypeptide having a deduced amino acid sequence of Table 1 [SEQ ID NO:2 or 4]. Variants that are fragments of the polynucleotides of the invention may be used to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding ribH variants, that have the amino acid sequence of ribH polypeptide of Table 1 [SEQ ID NO:2 or 4] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of ribH.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding ribH polypeptide having an amino acid

sequence set out in Table 1 [SEQ ID NO:2 or 4], and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical over its entire length to a polynucleotide encoding ribH polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of Table 1 [SEQ ID NO:1 or 3].

The invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. An example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein.

The invention also provides a polynucleotide consisting essentially of a polynucleotide sequence obtainable by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or a fragment thereof; and isolating said DNA sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers described elsewhere herein.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding ribH

and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the ribH gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

5 For example, the coding region of the ribH gene may be isolated by screening using a DNA sequence provided in Table 1 [SEQ ID NO: 1 or 3] to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

10 The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays.

Polynucleotides of the invention that are oligonucleotides derived from the sequences of Table 1 [SEQ ID NOS:1 or 2 or 3 or 4] may be used in the processes herein as described, but
15 preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that may encode a polypeptide that is the
20 mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino
25 acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

30 In addition to the standard A, G, C, T/U representations for nucleic acid bases, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA bases may appear at such a designated position in the DNA

or RNA sequence, except it is preferred that N is not a base that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

5 In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

10 **Vectors, host cells, expression**

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

15 For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

25 Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, enterococci *E. coli*, streptomyces and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells; and plant cells.

30 A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as

5 baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

10 For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

This invention is also related to the use of the ribH polynucleotides of the invention for use as diagnostic reagents. Detection of ribH in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the ribH gene may be detected at the nucleic acid level by a variety of techniques.

Nucleic acids for diagnosis may be obtained from an infected individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification technique prior to analysis. RNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of prokaryote present in an individual, may be made by an

analysis of the genotype of the prokaryote gene. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to labeled ribH polynucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in the electrophoretic mobility of the DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method. See, e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).


Cells carrying mutations or polymorphisms in the gene of the invention may also be detected at the DNA level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to a nucleic acid encoding ribH can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 2.

Table 2

Primers for amplification of ribH polynucleotides

<u>SEQ ID NO</u>	<u>PRIMER SEQUENCE</u>
5	5'-ATAGGTATTGTTGTAGCGAGATT-3'
6	5'-ATTTGGGCGATTCCTTTAGATAC-3'

The invention also includes primers of the formula:



wherein, at the 5' end of the molecule, X is hydrogen, and at the 3' end of the molecule, Y is hydrogen or a metal, R₁ and R₃ is any nucleic acid residue, m is an integer between 1 and 20 or zero, n is an integer between 1 and 20 or zero, and R₂ is a primer sequence of the invention, particularly a primer sequence selected from Table 2. In the polynucleotide formula above R₂ is oriented so that its 5' end residue is at the left, bound to R₁, and its 3' end residue is at the right, bound to R₃. Any stretch of nucleic acid residues denoted by either R group, where m and/or n is greater than 1, may be either a

heteropolymer or a homopolymer, preferably a heteropolymer being complementary to a region of a polynucleotide of Table 1. In a preferred embodiment m and/or n is an integer between 1 and 10.

The invention further provides these primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying ribH DNA
5 isolated from a sample derived from an individual. The primers may be used to amplify the gene isolated from an infected individual such that the gene may then be subject to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be detected and used to diagnose infection and to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial
10 infections, more preferably infections by *Streptococcus pneumoniae*, comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having a sequence of Table 1 [SEQ ID NO: 1 or 3]. Increased or decreased expression of ribH polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase
15 protection, Northern blotting and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of ribH protein compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a ribH protein, in a sample derived from a host are well-known to those of skill in the art. Such assay
20 methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Antibodies

The polypeptides of the invention or variants thereof, or cells expressing them can be used as an immunogen to produce antibodies immunospecific for such polypeptides. "Antibodies" as used
25 herein includes monoclonal and polyclonal antibodies, chimeric, single chain, simianized antibodies and humanized antibodies, as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

Antibodies generated against the polypeptides of the invention can be obtained by administering the polypeptides or epitope-bearing fragments, analogues or cells to an animal,
30 preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature*

256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice,
5 or other organisms such as other mammals, may be used to express humanized antibodies.

Alternatively phage display technology may be utilized to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-ribH or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10,
10 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing
15 the polypeptides to purify the polypeptides by affinity chromatography.

Thus, among others, antibodies against ribH- polypeptide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants that form a particular aspect of this invention. The term "antigenically equivalent
20 derivative" as used herein encompasses a polypeptide or its equivalent which will be specifically recognized by certain antibodies which, when raised to the protein or polypeptide according to the invention, interfere with the immediate physical interaction between pathogen and mammalian host. The term "immunologically equivalent derivative" as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise
25 antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

The polypeptide, such as an antigenically or immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be
30 associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically

or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized"; where the complementarity determining region(s) of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem.* 1989: 264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, 1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS USA* 1984:81,5849).

Antagonists and agonists - assays and molecules

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of ribH polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bacteriocidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising ribH polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a ribH agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the ribH polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product

from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of ribH polypeptide are most likely to be good antagonists. Molecules that bind well and increase the rate of product production from substrate are agonists. Detection of the rate or level of production of product from substrate may be enhanced by using a reporter system. Reporter systems that may be
5 useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in ribH polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for ribH antagonists is a competitive assay that combines ribH and a potential antagonist with ribH-binding molecules, recombinant ribH binding molecules,
10 natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. The ribH molecule can be labeled, such as by radioactivity or a colorimetric compound, such that the number of ribH molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

15 Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide or polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing ribH-induced activities, thereby preventing the action of ribH by
20 excluding ribH from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense
25 molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of ribH.

Each of the DNA sequences provided herein may be used in the discovery and
30 development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation

facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the initial physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block ribH protein-mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine *et al.*, *Infect. Immun.* 60:2211 (1992)); to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial ribH proteins that mediate tissue damage and; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

The antagonists and agonists of the invention may be employed, for instance, to inhibit and treat diseases.

Helicobacter pylori (herein *H. pylori*) bacteria infect the stomachs of over one-third of the world's population causing stomach cancer, ulcers, and gastritis (International Agency for Research on Cancer (1994) Schistosomes, Liver Flukes and *Helicobacter Pylori* (International Agency for Research on Cancer, Lyon, France; <http://www.uicc.ch/ecp/ecp2904.htm>). Moreover, the international Agency for Research on Cancer recently recognized a cause-and-effect relationship between *H. pylori* and gastric adenocarcinoma, classifying the bacterium as a Group I (definite) carcinogen. Preferred antimicrobial compounds of the invention (agonists and antagonists of ribH) found using screens provided by the invention, particularly broad-spectrum antibiotics, should be useful in the treatment of *H. pylori* infection. Such treatment should decrease the advent of *H. pylori*-induced cancers, such as gastrointestinal carcinoma. Such treatment should also cure gastric ulcers and gastritis.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with ribH, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Streptococcus pneumoniae* infection. Also provided are methods whereby

such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector to direct expression of ribH, or a fragment or a variant thereof, for expressing ribH, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual from disease, whether that disease is already established within the individual or not. One way of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid.

A further aspect of the invention relates to an immunological composition which, when introduced into an individual capable or having induced within it an immunological response, induces an immunological response in such individual to a ribH or protein coded therefrom, wherein the composition comprises a recombinant ribH or protein coded therefrom comprising DNA which codes for and expresses an antigen of said ribH or protein coded therefrom. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity or cellular immunity such as that arising from CTL or CD4+ T cells.

A ribH polypeptide or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Hemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilize the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof which have been shown to encode non-variable regions of

bacterial cell surface proteins in DNA constructs used in such genetic immunization experiments in animal models of infection with *Streptococcus pneumoniae* will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of
5 monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Streptococcus pneumoniae* infection, in mammals, particularly humans.

The polypeptide may be used as an antigen for vaccination of a host to produce
10 specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused, e.g., by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

15 The invention also includes a vaccine formulation which comprises an immunogenic recombinant protein of the invention together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection
20 solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials and may be stored in a freeze-dried condition requiring only the
25 addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

30 While the invention has been described with reference to certain ribH protein, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

Compositions, kits and administration

The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or their agonists or antagonists. The polypeptides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual

instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended
5 time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory peritoneal dialysis (CAPD) catheters.

The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device.
10 Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to prevent bacterial wound infections, especially *Streptococcus pneumoniae* wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibiotic prophylaxis before dental treatment that could produce a bacteremia.
15 Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to extend the use of the active agent as a replacement for prophylactic antibiotics in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins
20 exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibiotic prophylaxis.

Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1 µg/ml to 10mg/ml for bathing of wounds or indwelling devices.

25 A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to
30 suitable individuals.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety.

5 GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein.

"Disease(s)" means and disease caused by or related to infection by a bacteria, including otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and
10 endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid.

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art,
15 "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith,
20 D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine
25 identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990). The BLAST X program is publicly available
30 from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990). As an

illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant

method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be

appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide

may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1 Strain selection, Library Production and Sequencing

The polynucleotide having a DNA sequence given in Table 1 [SEQ ID NO:1 or 3] was obtained from a library of clones of chromosomal DNA of *Streptococcus pneumoniae* in *E. coli*. The sequencing data from two or more clones containing overlapping *Streptococcus pneumoniae* DNAs was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example: Methods 1 and 2 below.

Total cellular DNA is isolated from *Streptococcus pneumoniae* 0100993 according to standard procedures and size-fractionated by either of two methods.

Method 1

Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

Method 2

Total cellular DNA is partially hydrolyzed with a one or a combination of restriction enzymes appropriate to generate a series of fragments for cloning into library vectors (e.g., RsaI, PstI, AluI, BshI235I), and such fragments are size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the

vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E.coli* infected with the packaged library. The library is amplified by standard procedures.

5 Riboflavin Biosynthesis Operon

This ORF is part of an operon which encodes genes *ribG*, *ribB*, *ribA* and *ribH*. Gene *ribG* starts at nucleotide 1 and ends at nucleotide 1101. Gene *ribB* starts at nucleotide 1086 and ends at nucleotide 1721. Gene *ribA* starts at nucleotide 1711 and ends at nucleotide 2949. Gene *ribH* starts at nucleotide 2950 and ends at nucleotide 3417. This operon [SEQ ID NO:7] is as follows:

10 ATGAGCGATTCAAAATATATGAAATTAGCAATAAACTGGCACAAAAGGGGCTGGTTACGTCAATCCC
AATCCTATGGTTGGCGCAATTATTGTAAAAGATAATCACATTATCGGACAAGGTTATCATGAGTTTTTT
GGTGGCCACATGCTGAGAGAAATGCTCTTAAAACTGTAGAAAATCCCCTGTGCGAGCGACGCTTTAT
15 GTAACACTTGAACCCTGTTGTCACTTCGGGAAAACACCTCCCTGTATAGATGCTATAATCGATAGTGGT
ATTACAAGAGTAGTCATTGGAAGCCTAGACTGTAATCCTATTGTATCTGGAAAAGGAGTAAAGATACTT
GAGGAAAATAATCTTCAAGTTACTGTTGGAATTTTAGAAAATGAGTGTCTTAACCTAATAAAAAGTTTT
AGAAAGTATATTACCCAGCATGTACCCTATGTTTTTATGAAATATGCAATGTCAATGGATGGAAAATA
GCCACTAAAACAAATCAATCCAAATGGATTACTGAAGAAGAAGCAAGAAAGCATGTGCATCAGTTACGA
20 CACTATGTTAGTGCAATTATGGTGGGAGTCAATACTGTTATTCAAGACGATCCTTTGCTGACATGTAGA
TTGGAGGAAGGGAAAATCCTATCCGTATCATATGCGATACACATTTACGAACTCCTCTTACCTCTAAA
ATCGTAAAACAGCAAATGATATTAACCTTACATTGCCACTTCCTCTGAAGACAAAATAAAATGAAG
CTATATCAAAATCATGGCTGTGAAATACTTTCCATAAAGAAAAAGGCAATCATATAGACTTATCGAGT
TTAATGCAACATCTAGGAAACATGCAGATTGATAGCCTAGTTCTAGAGGGGGGAGTCTAATGAATTGG
25 AGTGCTTTGGAACAACAAATTGTTGATGAGCTGAAAATATATATTGCACCAAAAATTTTTGGAGGCAGT
GCCAAGTTTCCTGTGCGAGGTGAAGGCATTTCTTTGCCAAATGACGCTATTAGATTGAAACCTTATGCT
TTTTCTCAAATAGGAAATGACTATCTCATAGAAAGTGAAGTGATTTATCCATGTTACAGGAATAATTG
AAGAAATCGGAAAAGTTGAAAGAATACAGAAAGACTCTCGTAATTGTAACTATCAATTAAAGCCTCAA
AAATATTAACGGATATCCATTTAGGCGATAGTATAGCAGTAAATGGTATCTGTCTTACAGTTACTCATT
30 TCAATCATCAATCCTTTACAGTTGATGTAATGAATGAAACATGGAGTCGAACAGCTCTTACTCTATTAA
AACATGGAAGTGAGGTGAATCTAGAAAGAGCCITATCTGTCAACGGTCGACTTGGGGGTCACGTCGTTA
CAGGACACATTGATGGTACAGGAAAAATCTCGTCAATAAAAAAGATGATAATGCTGTATGGTATCAA
TCAACACACAAAAGAAATTTTAGATTTAATAGTTGAAAAGGATCTATTACAATTGACGGCATTAGTC
TGACTGTGCTAAAGTCTCCAAAGTAAACTTTTCAGTATCTGTTATCCCTCATACCTTGAAACAAACCA
35 TTCTTAAGAGTAAACAAGTCGGGAGTACAGTAAATCTTGAAATGATATCTTAGGTAATATGTGCAAA

AACTGATGGATAACTCTCCAAAATCAGAAATATCGAAGGAAGTATTATATCAAAATGGATTTTAGCAGA
AAGGATAATCAGTCAATGGAATATCGAAAAATGACAAGAAGCATTGAGGAAGCATTGCAGAAGGGACGA
CTTGTTCTTGTTATAGACGACAAGGATAGAGAAAATGAAGGAGACTTAATTTGTTCTGCACAAGCAGCT
ACAACAGAAAATGTTAATTTTATGGCTACTTATGCCAAAGGATTAATTTGTATGCCTATGAGCGAAAGT
5 TTAGCTAATCAATTAATGCTTTTACCTATGGTTGAAAACAATACAGATAATCATAAGACTGCTTTTACA
GTTTCAATTGATTATAAAGAAACGACCACAGGTATTTCTGCCGAGGAAAGAGGACTGACCGCACGTATG
TGTGTAGCTGAAGATATAACACCCTCTGATTTTCGCAGGCCAGGACACATGTTTCCTTTAATTGCAAAA
AAAGGTGGTGTCTAGAAAAGAAATGGACACACAGAAGCAACTGTTGATTTATTAATAATTAGCTGGACTA
AAAGAGTGTGGCCTATGTTGTGAAATAATGAATCATGATGGCAAAATGATGAGAACAGATGATTTAATT
10 CAGTTCTCGAAGAAACACAACATTCCTACTAATTACCATCAAAGAATTACAAGAATATAGAAAAGTATAT
GATCAGCTGGTAGAACGAGTTTCAACTGTCAATATGCCTACTAGATACGGTAATTTCAAAGCAATTAGC
TATATAGATAAACTAAATGGGGAACATCATCTTGCTCTTATTATGGGAAACATAGAGGATGAAGCCAAT
GTATTATGTCGGGTCCACTCCGAATGTTTAAACAGGAGATGTTTTAGGCTCTTTACGTTGCGATTGTGGA
CAGCAATTCGATAAAGCTATGAAAATGATTGTTGAGAATGGTTCGGGTGTCTTACTTTACTTGCGACAG
15 GAGGGACAAGGAATTGGACTTATCAATAAATTTAAAGCCTATCATTTACAAAATCAAGGCATGGATACG
CTTGATGCCAATCTTGCAATTAGGCTTTGAAGGTGATTTAAGAGAATATCATATTGGAGCACAAATGCTT
AAAGATCTGGGACTTCAGTCACTTCATTTACTGACAAATAATCCTGACAAGGTTGAACAGTTAGAAAAA
TATGGAATTACCATTTCCAGTAGAATATCAATCGAAATAGAAGCCAATCCTTACGATAGTTTTTATTTA
GAAACAAAGAAAAATCGAATGGGTCACATTTTAAATATGGAGGAAAAATAAATGAACACTTATGAAGGT
20 AATTTAGTAGCAACAATATTAAATAGGTATTGTTGTAGCGAGATTTAATGAATTTATAACTTCAAAA
TTATTATCTGGAGCACTAGATAATCTCAAAGAGAGAATGTAAACGAGAAAGATATCGAGGTAGCCTGG
GTTCCAGGAGCTTTTGAAATACCACTGATTGCATCAAAAATGGCAAAAAGTAAAAAATATGATGCAATT
ATCTGCTTGGGAGCTGTCATTAGAGGGAATACAAGTCATTATGATTATGTATGTAGCGAGGTATCTAAA
GGAATCGCCCCAATCAGTTTAAATAGCGAAATTCCTGTTATGTTTGGTGTGCTAACGACAGATACAATT
25 GAACAAGCCATAGAACGAGCTGGTACTAAAGCAGGAAATAAGGGTTCTGAGTGTGCACAAGGAGCTATT
GAAATGGTCAACCTAATTCGTACATTAGACGCATAG

30

What is claimed is:

1. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
2. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by the *ribH* gene contained in the *Streptococcus pneumoniae* of the deposited strain.
3. An isolated polynucleotide comprising a polynucleotide encoding a polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
4. An isolated polynucleotide that is complementary to the polynucleotide of claim 1.
5. The polynucleotide of Claim 1 wherein the polynucleotide is DNA or RNA.
6. The polynucleotide of Claim 1 comprising the nucleic acid sequence set forth in SEQ ID NO:1.
7. The polynucleotide of Claim 1 comprising nucleotide 1 to the stop codon which begins at nucleotide number 466 set forth in SEQ ID NO:1.
8. The polynucleotide of Claim 1 which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.
9. A vector comprising the polynucleotide of Claim 1.
10. A host cell comprising the vector of Claim 9.
11. A process for producing a polypeptide comprising: expressing from the host cell of Claim 10 a polypeptide encoded by said DNA.
12. A process for producing a *ribH* polypeptide or fragment comprising culturing a host of claim 10 under conditions sufficient for the production of said polypeptide or fragment.
13. A polypeptide comprising an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID NO:2.
14. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.
15. An antibody against the polypeptide of claim 14.
16. An antagonist which inhibits the activity or expression of the polypeptide of claim 14.

17. A method for the treatment of an individual in need of ribH polypeptide comprising: administering to the individual a therapeutically effective amount of the polypeptide of claim 14.

18. A method for the treatment of an individual having need to inhibit ribH polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 14.

19. A process for diagnosing a disease related to expression or activity of the polypeptide of claim 14 in an individual comprising:

- (a) determining a nucleic acid sequence encoding said polypeptide, and/or
- (b) analyzing for the presence or amount of said polypeptide in a sample derived from the individual.

20. A method for identifying compounds which interact with and inhibit or activate an activity of the polypeptide of claim 14 comprising:

contacting a composition comprising the polypeptide with the compound to be screened under conditions to permit interaction between the compound and the polypeptide to assess the interaction of a compound, such interaction being associated with a second component capable of providing a detectable signal in response to the interaction of the polypeptide with the compound;

and determining whether the compound interacts with and activates or inhibits an activity of the polypeptide by detecting the presence or absence of a signal generated from the interaction of the compound with the polypeptide.

21. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with ribH polypeptide of claim 14, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said animal from disease.

22. A method of inducing immunological response in a mammal which comprises delivering a nucleic acid vector to direct expression of ribH polypeptide of claim 14, or fragment or a variant thereof, for expressing said ribH polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody and/ or T cell immune response to protect said animal from disease.

23. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO:4.

24. An isolated polynucleotide comprising a polynucleotide having at least a 70% identity to the polynucleotide sequence of SEQ ID NO:3.

SEQUENCE LISTING

<110> SmithKline Beecham Corporation

<120> ribH

<130> P50533-11

<150> 08/978,454

<151> 1997-11-25

<160> 7

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 468

<212> DNA

<213> Streptococcus pneumoniae

<400> 1

atgaacactt atgaaggtaa tttagtagca aacaatatta aaataggtat tgtttagcg
60
agatttaatg aatttataac ttcaaaatta ttatctggag cactagataa tctcaaaaga
120
gagaatgtaa acgagaaaga tatcgaggta gcctgggttc caggagcttt tgaaatacca
180
ctgattgcat caaaaatggc aaaaagtaaa aaatatgatg caattatctg cttgggagct
240
gtcattagag ggaatacaag tcattatgat tatgtatgta gcgaggtatc taaaggaatc
300
gcccaaatca gtttaaatag cgaaattcct gttatgtttg gtgtgctaac gacagataca
360
attgaacaag ccatagaacg agctggtact aaagcaggaa ataagggttc tgagtgtgca
420
caaggagcta ttgaaatggt caacctaat cgtacattag acgcatag
468

<210> 2

<211> 155

<212> PRT

<213> Streptococcus pneumoniae

<400> 2

Met Asn Thr Tyr Glu Gly Asn Leu Val Ala Asn Asn Ile Lys Ile Gly
 1 5 10 15
 Ile Val Val Ala Arg Phe Asn Glu Phe Ile Thr Ser Lys Leu Leu Ser
 20 25 30
 Gly Ala Leu Asp Asn Leu Lys Arg Glu Asn Val Asn Glu Lys Asp Ile
 35 40 45
 Glu Val Ala Trp Val Pro Gly Ala Phe Glu Ile Pro Leu Ile Ala Ser
 50 55 60
 Lys Met Ala Lys Ser Lys Lys Tyr Asp Ala Ile Ile Cys Leu Gly Ala
 65 70 75 80
 Val Ile Arg Gly Asn Thr Ser His Tyr Asp Tyr Val Cys Ser Glu Val
 85 90 95
 Ser Lys Gly Ile Ala Gln Ile Ser Leu Asn Ser Glu Ile Pro Val Met
 100 105 110
 Phe Gly Val Leu Thr Thr Asp Thr Ile Glu Gln Ala Ile Glu Arg Ala
 115 120 125
 Gly Thr Lys Ala Gly Asn Lys Gly Ser Glu Cys Ala Gln Gly Ala Ile
 130 135 140
 Glu Met Val Asn Leu Ile Arg Thr Leu Asp Ala
 145 150 155

<210> 3

<211> 383

<212> DNA

<213> Streptococcus pneumoniae

<400> 3

ctagtttgct ctatgatttt cacagagtat taaattgcga ttttgccaag tttctttatt
 60
 cgtctaaaag tagagtctgt: tctatgcgtc taatgtacga attaggttga ccatttcaat
 120
 agctccttgt gcacactcag aaccttact tcttgcttta gtaccagctc gttctatggc
 180
 ttgtttcaatt gtatctgtcg ttagcacacc aaacataaca ggaatttcgc tatttaaact
 240
 gatttgggag attcctttaa atacctcgcc acatacataa tcataatgac ttgtattccc
 300

tctaatagaca gctcccaagc aaataattgc atcatatattt ttactttttg ccatttttga
 360
 tgcaatcagt gggatatttca aaa
 383

<210> 4
 <211> 99
 <212> PRT
 <213> Streptococcus pneumoniae

<400> 4
 Leu Lys Tyr Pro Leu Ile Ala Ser Lys Met Ala Lys Ser Lys Lys Tyr
 1 5 10 15
 Asp Ala Ile Ile Cys Leu Gly Ala Val Ile Arg Gly Asn Thr Ser His
 20 25 30
 Tyr Asp Tyr Val Cys Ser Glu Val Phe Lys Gly Ile Ala Gln Ile Ser
 35 40 45
 Leu Asn Ser Glu Ile Pro Val Met Phe Gly Val Leu Thr Thr Asp Thr
 50 55 60
 Ile Glu Gln Ala Ile Glu Arg Ala Gly Thr Lys Ala Gly Asn Lys Gly
 65 70 75 80
 Ser Glu Cys Ala Gln Gly Ala Ile Glu Met Val Asn Leu Ile Arg Thr
 85 90 95
 Leu Asp Ala

<210> 5
 <211> 23
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 5
 atagggtattg ttgtagcgag att 23

<210> 6
 <211> 23
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 6
 atttgggcga ttcctttaga tac 23

<210> 7

<211> 3417

<212> DNA

<213> *Streptococcus pneumoniae*

<400> 7

atgagcgatt caaaatatat gaaattagca ataaaactgg cacaaaaagg ggctgggttac
60
gtcaatccca atcctatggg tggcgcaatt attgtaaaag ataatcacat tatcggacaa
120
ggttatcatg agttttttgg tggccacat gctgagagaa atgctcttaa aaactgtaga
180
aaatcccctg tgggagcgac gctttatgta acacttgaac cctgttgtca cttcgggaaa
240
acacctccct gtatagatgc tataatcgat agtgggtatta caagagtagt cattggaagc
300
ctagactgta atcctattgt atctggaaaa ggagtaaaga tacttgagga aaataatctt
360
caagttactg ttggaatttt agaaaatgag tgtcttaact taataaaaag ttttagaaa
420
tatattacc agcatgtacc ctatgttttt atgaaatatg caatgtcaat ggatggaaaa
480
atagccacta aaacaaatca atccaaatgg attactgaag aagaagcaag aaagcatgtg
540
catcagttac gacactatgt tagtgcaatt atgggtgggag tcaatactgt tattcaagac
600
gaccccttgc tgacatgtag attggaggaa gggaaaaatc ctatccgtat catatgcat
660
acacatttac gaactcctct tacctctaaa atcgtaaaaa cagcaaatga tattaaaact
720
tacattgcca ctccctctga agacaaaaat aaaatgaagc tatatcaaaa tcatggctgt
780
gaaatacttt ccataaagaa aaaaggcaat catatagact tatcgagttt aatgcaacat
840
ctaggaaaca tgcagattga tagcctagtt ctagaggggg gcagtctaata gaattggagt
900
gctttggaac aacaaattgt tgatgagctg aaaatatata ttgcacaaa aatttttggg
960
ggcagtcca agtttcctgt cggaggtgaa ggcatttctt tgccaaatga cgctattaga
1020

ttgaaacctt atgctttttc tcaaataagga aatgactatc tcatagaaag tgaagtgatt
1080
tatccatggt cacaggaata attgaagaaa tcggaaaagt tgaaagaata cagaaagact
1140
ctcgttaattg taaactatca attaaagcct caaaaatatt aacggatatc catttaggcg
1200
atagtatagc agtaaatggt atctgtctta cagttactca tttcaatcat caatccttta
1260
cagttgatgt aatgaatgaa acatggagtc gaacagctct tactctatta aaacatggaa
1320
gtgaggtgaa tctagaaaga gccttatctg tcaacggtcg acttgggggt cacgtcggtta
1380
caggacacat tgatggtaca ggaaaaatct cgtcaataaa aaaagatgat aatgctgtat
1440
ggtatcaaat caacacacaa aaagaaatct tagatttaat agttgaaaaa ggatctatta
1500
caattgacgg cattagtctg actgtcgcta aagtctccaa agtaaacctt tcagtatctg
1560
ttatccctca taccttgaaa caaaccttc ttaagagtaa acaagtcggg agtacagtaa
1620
atcttgaaaa tgatatctta ggtaaatatg tgcaaaaact gatggataac tctccaaaat
1680
cagaaatatc gaaggaacta ttatatcaaa atggatttta gcagaaagga taatcagtc
1740
atggaatatc gaaaaatgac aagaagcatt gaggaagcat tgcagaaggg acgacttggt
1800
cttggttatag acgacaagga tagagaaaat gaaggagact taatttggtc tgcacaagca
1860
gctacaacag aaaatgttaa ttttatggct acttatgcc aaggattaat ttgtatgcct
1920
atgagcgaaa gtttagctaa tcaattaatg ctttcaccta tggttgaaaa caatacagat
1980
aatcataaga ctgcttttac agtttcaatt gattataaag aaacgaccac aggtatttct
2040
gccgaggaaa gaggactgac cgcacgtatg tgcgtagctg aagatataac accctctgat
2100
tttcgcaggc caggacacat gtttccttta attgcaaaaa aagggtggtg tctagaaaga
2160
aatggacaca cagaagcaac tgttgattta ttaaaattag ctggactaaa agagtgtggc
2220

ctatgttggtg aaataatgaa tcatgatggc aaaatgatga gaacagatga ttttaattcag
2280
ttctcgaaga aacacaacat tccactaatt accatcaaag aattacaaga atatagaaaa
2340
gtatatgatc agctggtaga acgagtttca actgtcaata tgctactag atacggtaat
2400
ttcaaagcaa ttagctatat agataaacta aatggggaac atcatcttgc tcttattatg
2460
ggaaacatag aggatgaagc caatgtatta tgtcgggtcc actccgaatg tttaacagga
2520
gatgttttag gctctttacg ttgcgattgt ggacagcaat tcgataaagc tatgaaaatg
2580
attgttgaga atggttcggg tgtcttactt tacttgcgac aggagggaca aggaattgga
2640
cttatcaata aattaaaagc ctatcattta caaaatcaag gcatggatac gcttgatgcc
2700
aatcttgcac taggctttga aggtgattta agagaatatc atattggagc acaaatgctt
2760
aaagatctgg gacttcagtc acttcattta ctgacaaata atcctgacaa ggttgaacag
2820
ttagaaaaat atggaattac catttccagt agaatatcaa tcgaaataga agccaatcct
2880
tacgatagtt tttatttaga aacaaagaaa aatcgaatgg gtcacatttt aaatatggag
2940
gaaaaataaa tgaacactta tgaaggtaat ttagtagcaa acaatattaa aataggtatt
3000
gttgtagcga gatttaatga atttataact tcaaaattat tatctggagc actagataat
3060
ctcaaaagag agaatgtaaa cgagaaagat atcgaggtag cctggggtcc aggagctttt
3120
gaaataccac tgattgcac aaaaatggca aaaagtaaaa aatatgatgc aattatctgc
3180
ttgggagctg tcattagagg gaatacaagt cattatgatt atgtatgtag cgaggtatct
3240
aaaggaatcg cccaaatcag tttaaatagc gaaattcctg ttatgtttgg tgtgctaacg
3300
acagatacaa ttgaacaagc catagaacga gctggtacta aagcaggaaa taagggttct
3360
gagtgtgcac aaggagctat tgaaatggc aacctaattc gtacattaga cgcatag
3417

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/25073

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 424/94.5, 244.1; 435/6, 193, 252.3, 325, 320.1; 536/23.2, 23.7

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/94.5, 244.1; 435/6, 193, 252.3, 325, 320.1; 536/23.2, 23.7

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS, MEDLINE, EMBASE

search terms: riboflavin; 6,7-dimethyl-8-ribityllumazine synthase; ribH; streptococcus pneumoniae

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FULLER, T.E. et al. Characterization of Actinobacillus pleuropneumoniae Riboflavin Biosynthesis Genes. Journal of Bacteriology. December 1995, Vol. 177, No. 24, pages 7265-7270, especially figure 3.	1-14, 23, 24



Further documents are listed in the continuation of Box C.



See patent family annex.

Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	*Z* document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

14 JANUARY 1999

Date of mailing of the international search report

01 MAR 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DIAN JACOBSON

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/25073

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

A61K 38/43, 38/45, 39/02; C07K 16/12, 16/40; C12N 1/21, 5/10, 9/10, 15/54, 15/63